## Where is the Origin of Human Consciousness?

## **Brain Mapping in Health and Disease**

By Jane Yip, PhD

317-503-1296

Indiana Brain Mapping www.IndianaBrainMapping.com


Autism Parent Care www.autismparentcare.com


#### Abstract


(1) Title: Brain mapping in health and disease

With the advent of neuroimaging in the last two decades, we have witnessed a quantum leap in the understanding of the brain. What was thought as idiosyncrasies of behavior is not only being taken more seriously but is being linked to certain loci in the brain. Just like discoveries of the microscope and telescope expanded our understanding, brain imaging has and will continue to challenge us into the question of consciousness and ultimately behavior. Technologies of neuroimaging has long been inaccessible for private clinicians as the NRI requires high cost for purchasing and maintenance. In the past 15 years, thanks to the help of computational science, brain imaging methods has been miniaturized into portable versions allowing much easier access and provide literally a map of the brain. The implication of such a trend will be discussed in this presentation along with its applications. Some results from case studies from individuals with brain conditions such as autism, ADHD, depression, addiction, learning disabilities, dyslexia and early onset Alzheimer's will be discussed along with strategies for intervention based on their brain mapping results.









## What makes the human brain unique?

Expansion of frontal lobe?


Emotion? Empathy

Language? Chimp language Bee language Birdsong

Blindsight NCC Binocular rivalry
Self-awareness?
Mirror test

## How do we study the human brain?







1848 Cavendish, VT


1953 William Scoville


# How is a brain put together?

"Ontogeny recapitulates phylogeny" - Ernst Haeckel 1866










# Anatomy: cortex






# Anatomy: cortex










# Animal models of cognition: bird song



# The New Era of Brain Imaging (1973 - )




Fig. 1- Comparative spatial and temporal resolution of different neuroimaging methods. The y-axis is the log10 of space and the x-axis is the log10 of time. The nested dynamics of the microscopic and mesoscopic levels being within the macroscopic level is illustrated in figure one. qEEG spatial resolution ranges from about 7 mm3 to 6 cm3 and temporal resolution less than 1 millisecond with the ability to measure events over a 24 hour period of time.

## **fMRI** (1990 - )

Functional magnetic resonance imaging, or fMRI, is a technique for measuring brain activity. It works by detecting the changes in blood oxygenation and flow that occur in response to neural activity










### Magnetoencephalography (MEG)

(2000 - )

Mapping brain activity by recording magnetic fields produced by electrical currents occurring naturally in the brain, using very sensitive








MEG is a record of magnetic fields, measured outside the head, produced by electrical activity within the brain.

### **Single-Photon Emission Computed Tomography (SPECT)**

(2000-)

A gamma-emitting radioisotope (called radionuclide) is injected into the subject's bloodstream, This allows the radioisotope to be carried and bound to a place of interest in the body, which then (due to the gamma-emission of the isotope) allows the ligand concentration to be seen by a gamma-camera.







Alzheimer's disease


## Electroencephalography (EGG) (1950 - )


Multi-frequency oscillations for scaling up or down in brain dynamics: The macro-, meso- and







Taken together, what have we learnt from brain imaging technology?











# Default Mode Network (DMN) Making sense of the Map: interpretation Making sense of the Map: interpretation Figure 1. A, B, A comparison of activity decreases during the performance of goal-directed tasks in the DMN (A) and spontaneous BOLD fluctuations (B) within the same network in the resting state derived from regions of interest placed in the posterioringuistel/precuneus (yellow arrow) or the medial perforntal cortex (orange arrow). The remarkable degree of coherence between activities obtained from these two distant loot within the same system is evident on inspection of these time-activity curves. The degree of system-wide coherence within the DNNs is shown in C when all voxels in the brain are examined for their posterior in function of the posterior original preformation (and arrow in A). Such a well connections. This figure was adapted with permissionfrom M. E. Raichle (2009).














## Live brain recording, a demo

Calling for a brave volunteer to demonstrate his/her thoughts live ....